Aspirus Langlade Hospital RN - Competency Assessment (Alteplase) Administration Guideline

	1 0	\ 1 /	
Name:		_Employee Number:	_ Job Class:

Competency:	Steps of Process:	Completed	Validators Signature & Date
Technical Domain:			
Demonstrates the ability to correctly mix and infuse Alteplase for Code Stroke patients who meet criteria. (Includes return demonstration)	 ☐ Get Alteplase kit from medication room. Contents include everything required for Alteplase infusion for Code Stroke patient. ☐ Remove cap from water vial 	□ pass □ fail	
	☐ Unwrap spike and maintain sterile condition		
Reason for Evaluation N = New Procedure/Equipment X H = High Risk X L = Low Volume X P = Problem Prone M = Mandatory D = Other Age Group N = Neonate D C = Child A = Adolescent X AD = Adult X G = Geriatric L = All	☐ Holding the vial of water vial in the upright position, spike the vial		
	☐ Remove the cap from the powdered Alteplase vial		
	☐ Hold the water vial firmly in the upright position		
	☐ With the other hand, hold the Alteplase vial upside down and lower Alteplase vial spike onto the water vial port		
	☐ With both hands, invert both vials together so the water will be on top and flow into the Alteplase vial		
	☐ Once all the water is in the Alteplase vial, grasp flanges of spike and decannulate the Alteplase vial		
	☐ Mix solution gently by rolling the Alteplase between your hands. DO NOT SHAKE. If there are bubbles in the vial, you will need to discard as it takes a long time for the bubbles to dissipate.		
	☐ Puncture Alteplase vial with IV tubing spike and open the vent on the side. Prime tubing with Alteplase.		
	☐ Utilizing the Alteplase dosing card, and after calculating the patient's weight in kilograms: locate the total Alteplase dose, Discard quantity, bolus dose and infusion dose.		

Aspirus Langlade Hospital RN - Competency Assessment (Alteplase) Administration Guideline

Name:	Emp	oloyee Number:	Job Class:
	will ALWAYS be disca mg) • Bolus dose – over 1 min • Infusion Dose = 58.9 kg	ml 6 ml (includes the 10 ml that arded as the maximum dose is 9 nute = 6.5 ml	0
	pump with above information.		
	exchange it for a 0.9 Normal Sakit in the medication room) and AS THE ALTEPLASE for one	☐ Once the infusion is complete, discard the Alteplase vial and exchange it for a 0.9 Normal Saline, 50 ml bag (in the Alteplase kit in the medication room) and infuse AT THE SAME RATE AS THE ALTEPLASE for one (1) hour	
	15 minutes during the infusion. NOTE – If you do not infuse to	checks must be completed every the Alteplase for some reason, cility is provided a new Alteplase	<u>se</u>
Action Plan with Remediation	Date and Level of Performance:	☐ deemed competent	□ not yet deemed competent
Job Class	Work Area		<u> </u>
Employee Signature	Date	Director, Coordinator or Validator Signature Date	